

The Secretary  
Senate Community Affairs References Committee  
Parliament House  
Canberra

Dear Secretary and Committee,

### **Inquiry into workplace exposure to toxic dust**

In my submission to the Inquiry, I wish to focus the Committee's attention on the **Alumina industry** where workers and surrounding communities have been exposed to numerous toxic elements contained in:

- bauxite dust
- caustic chemicals used to process bauxite into alumina
- the alumina itself
- dusts produced from liquor burning and oxalate destruction
- the residue or red mud.

I have summarized the hazards under various headings below and would be pleased to provide further information at the public hearings if required.

This submission is part of my wider study of hazards in the alumina industry.

I believe the Committee should examine the health and cause of death of former alumina industry workers, particularly those employed in the early years from 1960 to the 1980s as these workers can be expected to have developed more occupational disease.

Yours sincerely,

Geoff Pain Ph.D. B.Sc.(Hons.)

PO Box 1578  
Canning Vale  
WA 6970

## Toxic Dust Hazards in the Alumina Industry

Workers are exposed to heavy airborne dust loads from bauxite loading and grinding, caustic dusts inside washers, thickeners and sand traps, dusts from calciners, the liquor burner, oxalate kiln, alumina and hydrate dusts in the bagging department and residue (red mud) when driving to the red mud lakes.

Over 50% of red mud particles are less than 10.0  $\mu\text{m}$  [Glenister 1990].

Alumina producers accumulate red mud waste and are not permitted to dump it in the sea, one reason being that it leads to agglutination of fish gill tissues [Dethelsen, Sigmond]. It must be kept wet at all times and producers have emergency crews with tankers if sprinkler systems fail.

Worksafe time-weighted average limits are 10  $\text{mg}/\text{m}^3$  for inhalable and 5  $\text{mg}/\text{m}^3$  for respirable dusts. The NEPM for PM10 is 0.05  $\text{mg}/\text{m}^3$  for a one day average and the NHMRC interim indoor air quality goal is 0.09  $\text{mg}/\text{m}^3$ .

Alcoa Kwinana reported 410,000 kg of particulate matter below 10.0  $\mu\text{m}$  emitted to air annually in 1998-9 [NPI Kwinana]. This is 1200 kg per day. It would require 120,000,000  $\text{m}^3$  of air to dilute it to 10  $\text{mg}/\text{m}^3$ . This equates to a square kilometer to a height of 120 metres. Clearly on no wind days this level would be exceeded.

In 1997 it was **assumed in the absence of measurements** that dust emissions from the liquor burner at Kwinana would be 250  $\text{mg}/\text{m}^3$  [Galton-Fenzi].

The liquor burner at Kwinana emitted dust at the level of 39  $\text{mg}/\text{m}^3$  in December 2001.

Animal studies of dust load have been made.

Male rats were injected into their trachea with a suspension of dust of ore mine dead rock of high quartz content and mixed composition (particle size: 0-60 micrometers) [Tatrai].

Three and six months after treatment, foreign body granulation and after twelve months a non-fibrotic diffuse lung disease was seen in the animals.

After twelve months of exposure, non-specific, mature sinus histiocytosis and small focal epitheloid cell reaction were found to develop in the cervical, lung-hilar and retroperitoneal lymph nodes. Comparison of the histopathological changes with those seen in experimental silicosis induced by DQ12 quartz showed that the hazardous effect of industrial dusts is not identical with their fibrogenic effect.

At least one study has been made on particle deposition efficiency of artificial aerosols in the human maxillary sinus [Hyo N]. Deposition efficiencies in the human maxillary sinus were found to be 3%, for particles of 3-10 microns in diameter.

It has been found that 4-fold or greater risk increases of cancer of the nasal cavity and sinuses were associated with a history of chronic nasal diseases, including those occurring 10 or more years prior to cancer diagnosis [Zheng et al].

In 2003 Alcoa lost its self-policing rights for dust after it was disclosed that an employee had **falsified dust level recordings** on a database and Alcoa was fined for breaching dust emissions from its mud lakes [Southwell y].

Measurements of dust during manual descaling found average levels of 2.9 with a geometric standard deviation of 3.2 mg/m<sup>3</sup>

Other measurements comparing various sites at the Alcoa plant showed manual descaling with the high dust exposure of 7.9 (single measurement) and maximum levels of 1.5 for a residue operator, 5.6 for calciner operator/train loading and 8.1 reclaimer for operator. One sample measured 17.0 mg/m<sup>3</sup> for a calciner operator and this was rejected as excessive sample for further analysis.

The mass median aerodynamic diameter (MMAD, defined as the equivalent aerodynamic diameter such that 50% of the mass of the sample is associated with smaller particles) was found [Terry 1998] from 8 valid tests (from 11 measurements) to be 22.5 µm with a geometric standard deviation of 1.6 mg/m<sup>3</sup>

Another measurement during manual descaling found the MMAD to be 18.8 µm.

Although much is made of small dust particles' health risks due to their potential to travel far into the lung, risks associated with larger particles should not be discounted.

### ***Underestimation of Dust Loads***

Measurement of dusts can be difficult and it was shown in 1998 that measurements performed on Western Australian mine sites, including Alcoa sites, prior to that study were likely to have **underestimated dust loads by a factor of 2.4 to 3.4** [Terry 1998].

The authors emphasized that this would affect epidemiological investigations and compliance with statutory exposure standards.

All **Alcoa measurements listed here have not been corrected** on the basis of the 1998 findings by Terry *et al.*

The study showed that the previously used 7-hole (7H) and closed face (CF) samplers should be replaced with the Institute of Occupational Medicine sampler.

Alcoa made 43 measurements of "total" dust in K56 Digestion Building 17 from January 1990 to December 1998, averaging 7 to a maximum of 119 mg/m<sup>3</sup> over the period with the occupational exposure limit of 10 mg/m<sup>3</sup> exceeded on 6 sampling occasions.

Alcoa made 8 measurements of "respirable" dust in K53 descalers from September 1979 to November 1993, showing a maximum of 31.1 and a mean of 6.3 mg/m<sup>3</sup> over the period. The OEL was exceeded in 3 samples.

Alcoa made 4 measurements of "respirable" dust in K53 descale from January 1996 to December 2000, showing a decline from 1 to 0.2 mg/m<sup>3</sup> over the period.

Alcoa made 23 measurements of respirable dust in K56 Digestion Building 15 from January 1990 to December 1998, averaging 0.59 to a maximum of 1.58 mg/m<sup>3</sup> over the period .

Alcoa made 28 measurements of respirable dust in K56 Digestion Building 16 from January 1990 to December 1998, averaging 0.47 to a maximum of 3.1 mg/m<sup>3</sup> over the period .

Alcoa made 27 measurements of respirable dust in K56 Digestion Building 17 from January 1990 to December 1998, averaging 0.38 to a maximum of 1.2 mg/m<sup>3</sup> over the period .

### ***Alumina a possible carcinogen***

One of the most prominent dusts at refineries is of course alumina (aluminium oxide).

Alumina is classified by the International Agency for Research on Cancer (IARC) in Group 3, i.e. cannot be classified as to carcinogenicity to humans.

Although so far not confirmed as a carcinogen in humans,  $\gamma$ -alumina, formed by heating alumina hydrate to 900 to 1000° C, has been found to be highly fibrogenic when injected into rat lungs, with carcinogenic potential.

### ***Lung Overload associated with Alumina and Bauxite***

In a Mines Department study of the relationship between respiratory disease and particle size inhaled by workers [Hewson and Terry 1996], there is an interesting section on alumina and bauxite dusts that refers to lung overload.

Particulates often designated PNOC (particulates not otherwise classified) by the ACGIH TLV committee might in fact be more hazardous than assumed.

In rats at lung dust loadings of  $1 \text{ mg g}^{-1}$  lung tissue, particle clearance is delayed, however at  $10 \text{ mg g}^{-1}$  particle pulmonary clearance virtually ceased, leading to the term **dust overloading**. It was found that macrophages containing about 150 particles become immobilized and may release enzymes and other substances causing fibrosis [Morrow 1988, Vincent 1989 Muhle 1990, Mastomatteo 1993].

It was found volume, not mass is critical to immobilization.

Hewson and Terry state that dusts previous considered nuisance dusts could induce cancer and that current exposure standards for respirable dusts could be an order of magnitude too large [Pritchard 1989].

The ACGIH proposed a respirable dust standard of  $3 \text{ mg m}^{-3}$  for PNOC compared with the Western Australian standard of  $5 \text{ mg m}^{-3}$

## **Arsenic**

Arsenic was detected in Kwinana refinery ambient air in March 2002.

Sampling was done at typical worker locations including "in the liquor burner plume".

Arsenic was detected in:-

|             |                                      |
|-------------|--------------------------------------|
| Building 30 | in heater steam plume                |
| Building 48 | Granivor overflow vent odour         |
| Guardhouse  | Outside in liquor burner plume       |
| Building 44 | Baro Flash tank PRV vent steam       |
| Building 25 | #6 Rodmill discharge top door        |
| Building 51 | Tank top walkway liquor burner plume |

The maximum arsenic level detected was  $0.15 \text{ } \mu\text{g/m}^3$ . It was confirmed that the liquor burner plume descends to ground level within the plant.

A study on the occupational etiology of sinonasal cancer in three cases, which had arisen among glass workers, included an adenocarcinoma, a melanoma, and a squamocellular carcinoma. Evidence was found for cancer induction by arsenic dust [Battista 1996].

Increased incidence of bladder cancer has been found at a sulfuric acid factory with airborne arsenic levels of  $11 \text{ micrograms/m}^3$  [Englander 1988].

Arsenic inhalation is known to be associated with skin cancer [Hewson Terry 1996].

Arsenic is present in mud thickener launder scale (26.5 ppm) and floor scale 12 ppm.

The USEPA has set a target of zero for arsenic in drinking water.

## ***Beryllium***

The Committee will no doubt receive many submissions on this element following the recent publicity concerning beryllium disease in Navy personnel arising from descaling of ship surfaces.

Beryllium is present in red mud thickener launder scale (0.2 ppm) and floor scale (0.1 ppm).

Beryllium is present in red mud waste at 0.7 ppm.

The US National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, considers workers as exposed if the airborne concentration of beryllium is 100 ng/m<sup>3</sup> [Henneberger].

This concentration will be present if red mud dust is present at 143 mg/m<sup>3</sup>

Beryllium is known to induce debilitating and often fatal disease at low chronic exposures and significant lung function improvement may be seen following cessation of beryllium exposure [Cullen 1987, 2004].

A useful test was devised for chronic beryllium disease involving the proliferative response of bronchoalveolar lymphocytes to beryllium [Cullen 1989]

Screening for chronic beryllium disease will be vital for alumina and aluminium workers [Cullen 2005].

Beryllium compounds are irritating to the respiratory system and can cause skin sensitization.

## ***Diesel particulates***

Work inside alumina refinery tanks operating diesel equipment is comparable to underground mining where the respirable fraction of dust is about 50% of the inspirable aerosol [Terry 1998].

The following table is a summary of literature on cancer induced by diesel particulates.

| Cancer           | Odds Ratio or relative risk  | Year | Reference                         |
|------------------|--|------|-----------------------------------|
| laryngeal cancer | OR 5 - 5.4   | 1994 | Goldberg                          |
| lung             | 70 -year unit risk<br>2.1-5.5 per 10,000 workers per<br>$\mu\text{g}/\text{m}^3$ | 2001 | Dawson<br>Alcoa 8.35              |
| lung             | RR 1.3 95% CI 1.3-1.4  | 2001 | Bofetta<br>Alcoa 8.34             |
| lung             | 1.7 95% CI 0.58-6.0<br>after 20 years exposure                                   | 1999 | Saverin<br>Alcoa 8.27             |
| lung             | RR 1.63 95% CI 1.14 – 2.33   | 2000 | Gustavsson<br>Alcoa 8.31          |
| lung             | RR 2.4 95% CI 1.3 – 4.5  |      | Gustavsson<br>Alcoa 8.6           |
| lung             | RR 1.44 95% CI 1.01-2.05<br>Adjusted for smoking                                 | 2000 | Larkin<br>Alcoa 8.32              |
| lung             | 2.31 95% CI 1.44-3.70<br>heavy equipment operators                               | 1999 | Bruske-<br>Hohlfeld<br>Alcoa 8.25 |
| lung             | OR 1.43 95% CI 1.23-1.67<br>Adjusted for asbestos and<br>smoking                 | 2000 | Bruske-<br>Hohlfeld<br>Alcoa 8.29 |
| lung             | RR 1.47 95% CI 1.29-1.67   |      | Lipsett<br>Alcoa 8.26             |
| lung             | OR 1.21 95% CI 0.78-2.02   | 1990 | Alcoa 8.5                         |
| Sinonasal        | SIR 1.16 95% CI 0.98 – 1.36  | 2001 | Bofetta<br>Alcoa 8.34             |

## ***Thorium***

Alumina refinery workers are exposed to dust containing up to 400 ppm Thorium. This is more than 100 times the thorium concentration of basalt and 40 times the average for continental upper crust.

Thorium is an  $\alpha$ -particle emitter. A series of unstable daughter decay products are formed, including radium and radon.

Thorium was found in the 1960s to cause maxillary sinus cancer [Goren]. An x-ray contrast medium called Thorotrast was previously injected and shown to remain in the sinus for years [Galati].

Squamous cell Carcinoma of the parotid gland was found in 2 cases following sialography with Thorotrast 28 and 45 years previously [Nielsen 1979]. Both cases were histologically established as carcinoma and the presence of Thorotrast in the tumours was confirmed by autohistoradiography.

It was suggested that the tumours may have developed from metaplastic ductal epithelium after many years of exposure to the alpha radiation from Thorotrast deposits in the gland.

### ***Estimation of risk from radiation exposure at Kwinana***

Risk from inhalation of radioactive material at Kwinana requires knowledge of breathing rate, assumed to average  $1.2 \text{ m}^3/\text{hr}$ , 2000 hours per year, average radon and inhaled dust concentrations and composition of the dusts.

Unfortunately the amount of data is limited.

Estimates of geometric mean dose of 5.0 – 7.8 milliSieverts (mSv) have been made for underground bauxite mines in Europe [Bernhard 1984 cited in Hewson 1991].

This level of exposure would result in up to 200 excess cancers per million person years based on average energy dose to the whole body [Selinger 1989], however these figures do not allow for specific tissue susceptibility and damage power of particular isotopes.

When considering isotopes that decay in series, the energy contributions of all daughters (or progeny) have to be added.

### ***Estimates of excess deaths from inhalation of radionuclides at Kwinana***

Tables of mortality from cancers induced by inhaled radionuclides are available [Eckerman 1998].

Considering the data for Thorium-232 and its progeny only, a figure of approximately 6 deaths per million workers is expected for either acute one-time or chronic inhalation of **one Becquerel**. Note this **excludes** contributions of Radon- 220, Polonium-212, Polonium 216 and Thallium-208.

The red mud dust at Kwinana contains 400 ppm Thorium. Using this figure we can calculate that Thorium alone contributes at least 1.5 Becquerel per gram activity to the dust. This is in agreement with limited published data of up to 26 Becquerel per gram total activity for bauxite residue [O'Connor].



If we conservatively assume 3 Becquerel per gram total activity for bauxite residue at Kwinana and a dust level of 10 mg/m<sup>3</sup> a worker exposed for 1920 hours per year and breathing 1.2 m<sup>3</sup> / hour would inhale 23 g/year or 69 Becquerel.

The cancer mortality rate is therefore 414 per million.

If the worker is breathing this amount for 20 years, the cancer mortality rate is therefore 8280 per million. As noted above the actual inhalation of radionuclides will vary for each worker.

To this figure must be added the excess deaths expected from external radiation at the plant.

## **Silica**

Silica makes up about 21% of red mud waste. The committee will no doubt receive much data on this hazard in other submissions.

Silica is usually associated with lung cancer [Bruske-Hohlfeld, IARC Silica].

Nasal cancer risk in foundries show odds ratios up to 5.9 and this could be at least partially due to airborne silica [Comba 1992].

Alcoa made 4 measurements of silica in K53 descale from January 1996 to December 2000. Concentrations ranged from 0.0005 to 0.04 mg/m<sup>3</sup>

Alcoa made 8 measurements of silica in K56 Building 17 digestion from January 1996 to December 1998. Concentrations ranged from 0.002 to 0.09 mg/m<sup>3</sup>

## **Sodium hydroxide (caustic soda) and Bayer liquor**

Workers at alumina plants are required to use recycled caustic waters to hose down items of equipment as directed. During this hosing, caustic mists and, in dry air, dusts are generated.

Prolonged exposure to high concentrations of caustic soda can cause ulceration of nasal passages and severe eye and skin injury.

Irreversible obstructive lung disease following chronic occupational exposure as well as after a one-time high-level exposure to sodium hydroxide is known [Hansen and Isager 1991, Rubin *et al* 1992].

In 1969 the carcinogenic potential of sodium hydroxide was known.

Damage to the esophagus by sodium hydroxide can induce cancer scarring after swallowing [Lansing 1969]. Malignancy may develop even years after the removal of such tumors in the remaining part of the gullet. Total oesophagectomy was suggested instead of bypass [Csikos].

The patients with scar cancer comprised 7.2% of the overall oesophageal carcinoma cases. The interval between the caustic burn and the diagnosis of scar carcinoma was found to be 46.1 years. It was 50.9 years in those patients who drank lye before the age of 12, but 14 years less when it happened in adulthood.

Sodium and potassium hydroxide induce skin cancer with a long latency period in mice [Ingram].

Caustic substances cause tissue destruction through liquefaction or coagulation reactions and the intensity of destruction depends on the type, concentration, time of contact and amount of the substance ingested [ACGIH].

Rats that inhaled unmeasured concentrations of sodium hydroxide aerosols for 30 minutes per day suffered pulmonary damage after 2.5 months. [Mamede]

Workers at Kwinana report numerous rat deaths.

In humans, burning/redness of the nose, throat and eyes is found on exposure to caustic mists with NaOH concentrations between 0.005 and 0.7 mg/m<sup>3</sup>.

Alcoa made 11 measurements of NaOH mists in K53 Descalers from September 1979 to November 1993, averaging 0.26 to a maximum of 0.45 mg/m<sup>3</sup> over the period.

Alcoa made 57 measurements of NaOH mists in K58 Precipitation Building 45 from November 1993 to May 2002, to a maximum of 2 mg/m<sup>3</sup> over the period.

Alcoa made 40 measurements of NaOH mists in K58 Precipitation Building 46 from November 1993 to May 2002, averaging 0.2 to a maximum of 2 mg/m<sup>3</sup> over the period.

Alcoa made 20 measurements of NaOH mists in K58 Precipitation Building 60 from November 1993 to May 2002, averaging 0.2 to a maximum of 15.4 mg/m<sup>3</sup> over the period. The occupational exposure limit was exceeded by a factor of nearly 8 times on one sampling occasion.

On August 9, 2002, Alcoa issued a "World Wide Health Protocol - Mandatory Requirement" for all personnel exposed to dust, welding fume or caustic fumes to fill in a questionnaire annually.

Hill Brothers Chemical Company, a major supplier of sodium hydroxide warns of cancer risk in its Material Safety Data Sheet [Hill Brothers].

Bayer Liquor, a mixture of sodium aluminate, hydroxide, sulfate and chloride has its own Alcoa MSDS. It says "Vapors and mists can cause severe irritation of respiratory tract and lung damage". Alcoa recommends use of a personal respirator if the caustic mist exceeds 2 mg /m<sup>3</sup>.

In 2003, environmental sampling for operation and maintenance workers found "unacceptable" caustic mist concentrations in calcination at KW 59 (building 50) and "significant" in **K58** (buildings 45 and 46) and K28.

On the basis of these measurements, the Occupational Hygienist recommended personnel must wear a minimum of Class P2 disposable respirator, precipitation tanks on caustic wash cycle must be barricaded with Class P2 respirators provided for people entering the area.

Caustic soda washing of precipitators was increased in 2000 to economize on contractor descaling costs.

## ***Sodium oxalate***

This compound is a waste product, sent to kiln for thermal destruction or stored at red mud lakes. Single exposure causes severe redness and swelling of mucous membranes [Merck].

A very rare disease, antrolithiasis, involving deposition of calcium oxalate nodules, is most commonly found in maxillary sinus [Nass Duce 2003, Mori 2000].

The possibility that inhaled oxalate dusts at Kwinana result in precipitation of calcium oxalate particles in the sinus and other organs, especially the kidneys, should not be discounted. The carcinogenic potential of oxalate solids in the sinus should be investigated.

The sodium oxalate cake produced at Wagerup contains 0.9 ppm thorium, 3 ppm uranium, 84 ppm gallium, 20 ppm arsenic, 3.7 ppm mercury, 0.6 ppm nickel, 17 ppm selenium, 62 ppm vanadium.

When sent to the liquor burner or oxalate kiln, these elements can be expected in the particulate emissions.

Analysis of particulates from the Kwinana oxalate kiln found 15% below 2.5 µm, and 43% below 10 µm [Stackair 1997].

## ***Strong acid mists including sulfuric acid***

Alumina workers are exposed to strong acids in "acid cleaning" tasks e.g. when removing scale from cooling plates or similar equipment.

Strong acids are carcinogens of the upper and lower respiratory tract [Soskolne, NIOSH NTP].

Alcoa made 11 measurements over a period of 9 years of sulfuric acid mist in K56 Digestion Building 17 (labeled "draft only") with concentrations ranging from 0.001 to 13 mg/m<sup>3</sup>

Thus on one occasion (date not recorded) a level 13 times the occupational exposure limit was found.

Alumina workers are also exposed to sulfuric acid as one of the constituents of diesel engine exhausts.

In 1995 the WA Mines Safety and Inspection Regulations specified a maximum sulphur content of 0.5% compared to the tougher USA regulations of 0.05% [Terry 1998].

A study concluded that Western Australian industry, were not using best practicable technology and suggested revision of Australian Standard 3640 and recommended immediate implementation of control measures to reduce diesel emissions [Terry 1998].

## ***Vanadium compounds***

A Western Australian study of minesite dusts identified respirable vanadium as a health concern [Terry 1998].

Red mud waste contains over 400 ppm vanadium.

Vanadate exposure of rats significantly decreased the pressor response to 1-noradrenaline, and increased the pressor response to sinus-carotid baroreceptor stimulation and the hypotensive responses to 1-isoprenaline and acetylcholine [Matera].

Vanadium pentoxide catalyzes formation of maleic anhydride from benzene in air.

In a Mines Department study of the relationship between respiratory disease and particle size inhaled by workers, vanadium dusts are said to produce **green tongue disease** and leave a metallic taste.

A number of workers complain of a metallic taste associated with alumina refinery emissions.

**Summary table of airborne dusts in the alumina industry with acute and chronic health limits and examples of measured levels from one company's reports, where available (PINJ = Pinjarra refinery data, LB = Liquor Burner, OX = Oxalate residue).**

| CHEMICAL  | ATSDR MRL                         |   | ALCOA<br>KWINANA<br>EMISSIONS In<br>$\mu\text{g}/\text{m}^3$ unless<br>otherwise marked |
|---|-----------------------------------|---|---|
| ARSENIC   | Provisional<br>Oral Acute<br>Chr. | 0.005 mg/ kg/ day<br><br>0.0003 mg/ kg/<br>day          | 48 ppm in mud<br>295 g/day PINJ<br>13.5 g/day LB  |
| BERYLLIUM                                       | Oral Chr.                         | 0.002 mg/ kg/ day                                       | 0.7 ppm in mud  |
| CADMIUM   | Oral Chr.                         | 0.0002 mg/ kg/<br>day                                   | 4.5 ppm in mud  |
| CHROMIUM(VI), AEROSOL<br>MISTS                  | Inh. Int.                         | 0.000005 mg/ m <sup>3</sup>                             |   |
| CHROMIUM(VI),<br>PARTICULATES                   | Inh. Int.                         | 0.001 mg/ m <sup>3</sup>                                | 314 ppm in mud  |
| COBALT  | Inh. Chr.<br>Oral Int.            | 0.0001 mg/ m <sup>3</sup><br>0.01 mg/ kg/ day           | 22 ppm in mud   |
| FLUORIDE, SODIUM                                | Oral Chr.                         | 0.05 mg/ kg/ day  | 1200 ppm in mud   |
| MANGANESE                                       | Inh. Chr.                         | 0.00004 mg/ m <sup>3</sup>                              | 300 ppm in mud<br>2125 g/day PINJ   |
| NICKEL COMPOUNDS                                | Inh. Int.<br><br>Chr.             | 0.0002 mg/ m <sup>3</sup><br>0.00009 mg/ m <sup>3</sup> | 65 g/day PINJ<br>5 ppm in mud   |
| PARTICULATES                                    |                                   |   | 200,000 LB<br>54,000 OX<br>271,296 g/day LB<br>528077 g/day<br>PINJ                     |
| SILICA  |                                   |   | 21% in mud  |
| SODIUM HYDROXIDE                                |                                   |   |   |
| STRONG ACID MISTS<br>INCLUDING SULFURIC<br>ACID |                                   |   |   |
| SELENIUM  | Oral Chr.                         | 0.005 mg/ kg/ day<br>3                                  | 120 LB<br>217 g/day PINJ<br>163 g/day LB<br>3 PPM in mud                                |
| SODIUM OXALATE                                  |                                   |   |   |
| VANADIUM  | Inh. Acute                        | 0.0002 mg/ m <sup>3</sup>                               |   |

## References used in the submission and other useful resources

Please note some are newspaper cuttings of unknown date

ABC Background Briefing "Alcoa's Bad Odour" Sunday 9 June 2002  
<http://www.abc.net.au/rn/talks/bbing/stories/s579605.htm>

Adolph F 2004 The Sunday Times February 29 Residents fear radioactivity

Adolph F 2004 The Sunday Times Air tests fail Alcoa probe

Adshead G 2004 The West Australian April 23 p2 Alcoa workers exposed to fumes  
-poisonous fumes hit Alcoa workers

Adshead G 2004 The West Australian April 24 p2 Bloody noses blamed on Alcoa

Alcoa 2002 Refinery Air Emissions Inventory Progress report 9 August 2002 2<sup>nd</sup> Edition

Amalfi C 2002a The West Australian Jan 29 p1 Forget It – lawyer questions Alcoa  
actions

Amalfi C 2002b The West Australian Jan 29 p6 Alcoa health case row : Plaintiff alleges  
pressure to drop court action

Amalfi C 2002 The West Australian August 31 p1 K58 death list – workers document  
health fears

Amalfi C 2002 The West Australian August 31 p11,12 Out of breath but not fight – lives  
lost and lungs ruined, say Alcoa staff at refinery

Amalfi C 2003 The West Australian March 15 p1 David takes on Alcoa, and wins

Amalfi C 2003 The West Australian March 15 p15 Former Alcoa worker wins illness  
compo – father of three claims double win against corporate Goliath

Amalfi C 2003 The West Australian March 18 p12 Judge accuses Alcoa – long-running  
case thrown out

Amalfi C 2003 The West Australian June 7 p Tests point to high cancer risk – fatal  
legacy hangs over Alcoa workers

Amalfi C 2003 The West Australian p Probe into deaths of Alcoa workers – concern over  
Kwinana refinery area called K58

Amalfi C 2003 The West Australian August 28 p7 Alcoa lobby wants new study – workers widow claims cancer risk whitewash

Amalfi C 2004 The West Australian p12 Alcoa cancer study urged

Amalfi C 2004 The West Australian April 29 p1 Alcoa faces fresh cancer inquiry

Amalfi C 2004 The West Australian April 29 p1 Other Alcoa sites may face health probe

Amalfi C 2004 The West Australian May 10 p 7 More cases included in Alcoa study

ATSDR 2005 Agency for Toxic Substances and Disease Registry, Minimum Risk Levels <http://www.atsdr.cdc.gov/mrls.html>

Bagley ST *et al* 2002 Impact of low-emission diesel engines on underground mine air quality Alcoa 8.37

**Battista G** *et al* 1996 Am J Ind Med. 30(1) 31-5. Art glassware and sinonasal cancer: report of three cases.

Benchmark Technology Services 2004 Peer review of Environ report on upgraded liquor burner.

**Bernhard S** *et al* 1985 J Radiol Prot 8 19-24 Occupational exposure to Rn in non-uranium mines

Bhargava SK 2000 TMS Annual meeting March 12-16 Nashville Organics removal from bauxite

Bigu J 1990 Electrical charge characteristics of long-lived radioactive dusts Health Physics 58 341-50

Bloom NS, Prestbo EM and Dobbs CL Geochemistry of mercury in the Bayer process

Bofetta P *et al* 2001 Cancer Causes Control 12 365-74 Occupational exposure to diesel engine emissions and risk of cancer in Swedish men and women Alcoa 8.34

Brinton LA *et al* 1984 Am J Epidemiol 119(6) 896-906 A case-control study of cancers of the nasal cavity and paranasal sinuses

Bruske-Hohlfeld I 1999 Am J Ind Med 36 405-14 Lung cancer risk in male workers occupationally exposed to diesel motor emissions in Germany Alcoa 8.25

Bruske-Hohlfeld I 2000 Amer J Epidemiology 151(4) 384-95 Occupational Lung cancer risk for men in Germany: results from a pooled case-control study

Butler J 2002 The West Australian May 18 p4 More time for ill to sue – Cabinet plan will add to insurance cost, say Foss and employers

Butler J 2002 The West Australian May 18 p4 Time limit questioned on killer compo cases

Butler J 2002 The West Australian August 14 p Yarloop residents lodge Alcoa writs -action reserves right to sue over liquor burner

Calverley A 2004 The West Australian Activists want ban on Alcoa refinery

Calverley A 2004 The West Australian November 25 p6 Another spill at Alcoa's refinery

Calverley A 2004 The West Australian November 26 p10 Alcoa dismisses critics' call to close Kwinana refinery

Colville A 2004 Community Weekend Courier September 24 p1 Assessment ruling for burner cheered

**Comba P et al** 1992 Br J Ind Med 49(3) 193-6 Cancer of the nose and paranasal sinuses in the metal industry: a case-control study.

**Comba P et al** 1992 b Am J Ind Med 22(4) 511-20 A case-control study of the nose and paranasal sinus and occupational exposures

**Comba P and Belli S** 1992 Ann Ist Super Nata 28(1) 121-32 Etiological epidemiology of the nasal cavities and paranasal sinuses

Cornet A *et al* Sem Hop 1976;52:1999-2003 Caustic burn and adenocarcinoma of the esophagus.

CSIRO Atmospheric Research 2004 Wagerup Air Quality Review

**Csikos M et al** 1985 Langenbecks Arch Chir. 365(4) 231-8. Late malignant transformation of chronic corrosive oesophageal strictures.

**Cullen MR et al** 1987 Am Rev Respir Dis. Jan;135(1):201-8. Chronic beryllium disease in a precious metal refinery. Clinical epidemiologic and immunologic evidence for continuing risk from exposure to low level beryllium fume.

**Cullen M R et al** Ann Intern Med. 1988 May;108(5):687-93. Erratum in: Ann Intern Med 1989 110(8):672. Proliferative response of bronchoalveolar lymphocytes to beryllium. A test for chronic beryllium disease.

**Cullen M R** 1996 Chest 109(3) Supplement 51S-54S Epidemiologic methods for the study of occupational asthma. Current problems and solutions.



**Cullen** M R 1990: Chest 98:196S-201S Clinical surveillance and management of occupational asthma.

**Cullen** MR *et al* Chest. 2004 126(6):1730-2. Variable response to long-term corticosteroid therapy in chronic beryllium disease.

**Cullen** MR 2005 Am J Respir Crit Care Med. Jan 1;171(1):3-4. Screening for chronic beryllium disease: one hurdle down, two to go. No abstract available.

Dauids PH Ned Tijdschr Geneeskd 2001 145(44) 2105-8 Consequences of caustic damage to the esophagus

Dawson S V and Alexeeff 2001 Risk Analysis 21(1) 1-18 Multi-stage model estimates of lung cancer risk from exposure to diesel exhaust based on a US railroad worker cohort

**Dethelsen** V and Rosenthal H 1973 Aquaculture 2 267-80 Problems with dumping of red mud in shallow waters.

Deitmer T 1989 Strahlenther Onkol. 165(6):444-6. Tumors of the inner nose and paranasal sinuses due to asbestos exposure?

Deitmer T, Borsch-Galetke E. HNO. 1993 41(7):352-5. Nose and paranasal sinus malignancies with reference to occupational medicine aspects. A case study

Deitmer T and Wiener C 1996 Acta Otolaryngol 116(5) 762-5 Is there an occupational etiology of inverted papilloma of the nose and sinuses? Ref 15 in Alcoa 8.39

Department of Health 2002 Report 4-52293 June 27 Report on cancer in Alcoa Workers

Diesel Emission Evaluation Program 2000 <http://www.deep.org/research.html>

Donoghue AM a 2001 Occup. Med. 51(2) 118-123 The design of hazard risk assessment matrices for ranking occupational health risks and their application in mining and minerals processing.

Donoghue AM b 2001 Occup. Med. 51(5) 354-356 A risk-based system to penalize and reward line management for occupational safety and health performance.

Donoghue AM c 2001 Occup. Med. 51(4) 266-71 The calculation of accident risks in fitness for work assessments: diseases that can cause sudden incapacity.

Donoghue AM 2003 email July 18 to Merran Smith re approaching new sinus cancer cases to seek link to Alcoa

**Eckerman** KF *et al* 1998 Oak Ridge National Laboratory USEPA Office of Radiation and Indoor Air, Federal Guidance Report No 13. Health risks from low-level environmental exposure to radionuclides. Radionuclide-specific lifetime radiogenic cancer risk coefficients for the US population based on age-dependent intake, dosimetry and risk models.

**Englander** V *et al* Int Arch Occup Environ Health. 1988;61(3):157-62. Mortality and cancer morbidity in workers exposed to sulphur dioxide in a sulphuric acid plant.

Environ 2004 Atmospheric emissions screening assessment upgraded Kwinana liquor burner

Environ 2004 Kwinana liquor burner emissions reduction project Works Approval Application

Eyer SL, Bhargava S, Sumich M, Tardio J, Akolekar D and Nyeboer H Light metals 2000 The Minerals, Metals and Materials Society 45-51 Removal of organics from Bayer liquor and wet oxidation

Flodin U *et al* 1981 Scand J Work Environ Health 7 169-78

Forster P 1998 23 January, to Ormonde N, Genoni S, Power G, Petrie H, Adamak G, Armanios C, Sonehouse L Union concerns with Kw liquor burning emissions

Fox AJ and Collier PF. Br J Prev Soc Med. 1976 30(4):225-30. Low mortality rates in industrial cohort studies due to selection for work and survival in the industry.

**Galati** L T, 2002, Malignant Tumors of the Sinuses  
<http://www.emedicine.com/ent/topic254.htm>

**Galton-Fenzi** B 1997 Liquor burning impact assessment (some health issues considered)  
4<sup>th</sup> June

Gauntlett K 2003 The West Australian January 21 p9 Alcoa in strife over altered dust readings

**Glenister** D 1990 Particle size distribution in red mud residue.

Gold LS *et al* 2003 Regulatory Toxicology and Pharmacology 37, 45-53 Comparison of cancer risk estimates based on a variety of risk assessment methodologies.

Goldberg M *et al* 1994 Cancer causes Control 5(1) 15-25 A 10-year incidence survey of respiratory cancer and a case-control study within a cohort of nickel mining and refining workers in New Caledonia

**Goren, AD** *et al* Oral surgery, oral medicine, oral pathology, oral radiology, & endodontics, 1980 49(3) 237-242 Clinical and radiobiologic features of thorotrast-induced carcinoma of the maxillary sinus - case-report

Granath N Alcoa sorry as another chemical leak sparks investigation

Gregory A 2002 The West Australian July 9 p12 Alcoa in denial over health claims: union

Groves J and Cain JR 2000 Ann Occup Hygiene 44(6) 435-47 A survey of exposure to diesel engine exhaust emissions in the workplace

Gustavsson P 2000 Am J Epidemiol 152(1) 32-41 Occupational exposure and lung cancer risk: A population-based case-referent study in Sweden

Gustavsson P *et al* 1990 Scand. J Work Environ Health 16 348-54 Lung cancer and exposure to diesel exhaust among bus garage workers

Halden T 2004 Worries over burner emissions

Hall T 2004 The Sunday Times December 5 Poison the pollies

**Hansen** KS and Isager H, J. Society Occup Med 1991 41(1) 45-6 Obstructive lung injury after treating wood with sodium hydroxide.

Harreus U *et al* 1999 Laryngorhinootologie 78(4) 176-81 Genotoxic effect on human mucous membrane biopsies of the upper aerodigestive tract

Hartley BM and Hewson GS 1993 Radiat Protect Aust 11 165-9 Summary of past and present radiation research into thorium (monazite) dosimetry

Henneberger PK J Occup Environ Hyg. 2004 1(10):648-59. Industries in the United States with airborne beryllium exposure and estimates of the number of current workers potentially exposed.

Hewitt S 2004 The West Australian June 30 p 4 Toxic overflow at Alcoa plant

Hewitt S 2004 The West Australian June 30 p 4 Mystery surrounds thyroid cancer figures

Hewitt S 2004 The West Australian p Another spill at Alcoa

**Hewson** GS *et al* 1991 Minerals and Energy Research Institute of Western Australia (MERIWA) Report No. 79 Preliminary study of radon in underground mines in Western Australia

**Hewson GS and Terry KW** 1996 Minerals and Energy Research Institute of Western Australia (MERIWA) Report No. 162 The relationship between respiratory disease and the particle size of dust inhaled by mine workers

**Hill Brothers** Chemical Company Material Safety Data Sheet.  
<http://www.hillbrothers.com/msds/sh-flake.htm>

**Hyo N** Rhinology. 1989 27(1):17-26. Particle deposition efficiency of therapeutic aerosols in the human maxillary sinus.

IARC Beryllium and Beryllium Compounds International Agency for Research on Cancer Tenth Report on Carcinogens

IARC Diesel Exhaust Particulates International Agency for Research on Cancer Tenth Report on Carcinogens

IARC Internally absorbed radionuclides International Agency for Research on Cancer Tenth Report on Carcinogens

IARC Radon International Agency for Research on Cancer Tenth Report on Carcinogens

**IARC Silica**, Crystalline (Respirable Size) International Agency for Research on Cancer Tenth Report on Carcinogens

IARC Strong Inorganic acid mists containing sulfuric acid International Agency for Research on Cancer Tenth Report on Carcinogens

IARC Thorium Dioxide International Agency for Research on Cancer Tenth Report on Carcinogens

**Ingram** AJ and Grasso 1991 Mutat Res 248(2) 333-40 Evidence for possible mechanisms of non-genotoxic carcinogenesis in mouse skin

Kerswell T 2004 Sound Telegraph September 29 p 1 Refinery protest rage – Edwards agrees to review

Langer AM *et al* Science. 1980 Jul 18;209(4454):420-2. Asbestos as a cofactor in carcinogenesis among nickel-processing workers.

**Lansing** PB 1969 *et al* Am J Surg 118, 108-111 Carcinoma of the esophagus at the site of lye stricture

Lareo AC *et al* 1992 Laryngoscope 102(4) 439-42

Larkin EK *et al* 2000 Am J Ind Med 38 399-409 Diesel exhaust exposure and lung cancer: adjustment for the effect of smoking in a retrospective cohort study Alcoa 8.32

Lipsett M and Campleman S Am J Public Health 89 1009-17

Luce D *et al* 2002 Cancer Causes Control 2002 Mar;13(2):147-57. Sinonasal cancer and occupational exposures: a pooled analysis of 12 case-control studies.

Luce D *et al* 1997 Am J Ind Med 32(3) 205-10 Sinonasal cancer and occupational exposure to wood dust

Magill P and Ruse B 2002 The West Australian March p Alcoa wins payout appeal

**Mamede** RC and de Mello Filho FV Sao Paulo Med J. 2001 Jan 4;119(1):10-5. Ingestion of caustic substances and its complications.

Marchman IW and Hewson GS 1995 Radiation doses and associated parameters in the Western Australian mineral sands industry Department of Minerals of Energy Perth WA

**Mastromatteo** E 1993 Is mining an unhealthy industry? Minesafe International Perth 1993

Matera MG Riv Eur Sci Med Farmacol. 1989 11(2):105-12. Prenatal and postnatal vanadium exposure in rats: effect on vasomotor reactivity development of pups.

Mayman J 2002 The Weekend Australian May 11-12 Inquirer 19-23

McCawley MA *et al* 2001 Applied Occupational and Environmental Hygiene 16 631-8 Ultrafine beryllium number concentration as a possible metric for Chronic Beryllium Disease Risk

McPharlin IR, Jeffery RC, Toussaint LF and Cooper M 1992 P, N and radionuclide retention and leaching in a Joel sand amended with red mud/gypsum

Melville Times 2004 May 18 p1 Inquiry win for widows

**Merck** Pty Ltd Material Safety Data Sheet (HazMat printout 14/3/2000) Sodium Oxalate

Miraudo N 2002 The Sunday Times p Cancer in WA up by 50 pc

Miraudo N 2002 The Sunday Times p Gift of Love – wife donates kidney to save husband

Miraudo N 2002 The Sunday Times p Alcoa expert links illness to refinery

**Mori** S *et al* J Otorhinolaryngol Relat Spec. 2000 62(6):335-7. Antrolithiasis in the frontal sinus.

Miraudo N 2002 The Sunday Times p Impartial study call for Alcoa

Miraudo N 2004 The Sunday Times p New bid to fight alumina

Morimoto K 1976 Jpn J Ind Health (Sangyo Igaku) 18 23-34 Analysis of the combined effects of benzene with radiation on chromosomes in cultured human leukocytes.

**Morrow** PE 1988 Fundam App Toxicology 10 369-84 Possible mechanisms to explain dust overloading of the lungs

**Muhle** H *et al* 1993 J Aerosol Med 3 s111-128 Dust overloading of lungs: Investigation of various materials, species, differences and irreversibility of effects

**Nass** D *et al* J Laryngol Otol. 2003 117(8):637-40. Antrolithiasis: a retrospective study.

National Cancer Institute 2003 cancer facts [http://cis.nci.nih.gov/fact/3\\_8.htm](http://cis.nci.nih.gov/fact/3_8.htm)

Nemery B 1990 Eur Respir J 3 202-19 Metal toxicity and the respiratory tract

**Nielsen** M *et al* Acta Otolaryngol. 1979;88(5-6):462-7. Carcinoma of the parotid gland following sialography with thorotrast. Report of two cases.

NIOSH NTP National Toxicology Program 10<sup>th</sup> report on Carcinogens: Strong Inorganic Acid Mists containing Sulfuric Acid

**NPI** National Pollutant Inventory Emission Report Alcoa Kwinana 1998-99

NPI Substance profile Selenium <http://www.npi.gov.au/database/substance-info/profiles/75/html>

Oberdorster G *et al* 1992 Environ Health Perspective 97 193-199 Role of the alveolar macrophage in lung injury: studies with ultrafine particles

**O'Connor** BH, Fox DR, Glenister DJ, Chesson BJ, Summers KJ and Smirk DD 1991 Radiological evaluation of proposed end-uses of solid residue from bauxite processing. Report to the Radiological Council of Western Australia

OEHHA 2005 Office of Environment Health Hazard Assessment, California Governor Arnold Schwarzenegger [www.oehha.gov/air](http://www.oehha.gov/air)

OEHHA 1999 Determination of Acute reference exposure levels for airborne toxicants Acute Toxicity Summary Sodium Hydroxide [www.oehha.gov/air/acute\\_rels/pdf/1310932A.pdf](http://www.oehha.gov/air/acute_rels/pdf/1310932A.pdf)

O'Leary C 2004 The West Australian October 5 p1 The postcodes that come with cancer

O'Leary C 2004 The West Australian October 5 p11 Cancer rates less in rural areas: report

O'Leary C 2004 The West Australian Exercise gives Kwinana residents a headache

ORNL 11 Oak Ridge National Laboratory Federal Guidance Report 11 look up tables  
<http://homer.hsr.ornl.gov/VLAB/FedGR11.html>

Ott MG *et al* 1977 J Occup Med 19 813-16 Mortality among employees chronically exposed to caustic dust.

Ott MG *et al* 1989 Am J Ind Med. 6(6):617-30. Assessment of exposure to chemicals in a complex work environment.

Ott MG *et al* 1989 Am J Ind Med. 16(6):631-43 Lymphatic and hematopoietic tissue cancer in a chemical manufacturing environment.

Paris C *et al* Rev Prat. 2004 54(15):1660-4. Lung cancer and occupational exposures (asbestos not considered)

Pederson TC and Siak J-S 1981 J Appl Toxicol 1 54-60 The role of nitroaromatic compounds in the direct-acting mutagenicity of diesel particle extracts.

Penn S 2004 Mine workers safety at risk, says activist

Pennells S 2004 The West Australian June 26 p1 \$10b resources plan in pipeline

Pennells S 2004 The West Australian June p 17 \$4m to hasten resource bids

PER Public Environmental Review Use of Bauxite Residue in the Peel-Harvey coastal plain catchment 1993

Pescud C 2004 Community Melville Times September 28 p1 Protest primed by cancer fears

Phaceas J 2004 The West Australian p Wagerup warning, Pinjarra approved

Phaceas J 2004 The West Australian p Alcoa kicks off plans for big expansion

Phaceas J 2004 The West Australian p Alcoa studies Guinea to speed up Wagerup talks

Phaceas J 2004 Approvals sought for Alcoa's Wagerup expansion

Phaceas J 2004 The West Australian p49 Production limits cloud Wagerup expansion

Pitsis S 2004 The Australian December 21 p Alcoa pleads guilty on refinery pollution

Power G 2002 personal communication to G Pain

**Pritchard** JN 1989 J Aerosol Sc. 20 1341-4 Dust overloading – a case for lowering the TLV of nuisance dusts?

Pryer W 2004 July 3 p40 Fears ignored on emissions: MLA

Reger RB and Morgan WKC 1993 Occupational Medicine 8 185-204 Respiratory cancers in mining

Rosenman KD and Stanbury M. Am J Ind Med. 1996 May;29(5):491-500. Risk of lung cancer among former chromium smelter workers.

**Rubin** AE *et al* Br J Ind Med 1992 49 213-14

Rudell B *et al* 1999 Occup Env Med 56 222-231 Efficiency of automotive cabin filters to reduce acute health effects of diesel exhaust in human subjects Alcoa 8.28

Ruse B 2002 The West Australian March 12 p 4 We will look after sick: Alcoa – Any health risks that are connected with the refinery at Wagerup are minor, according to Alcoa

Saverin R *et al* 1999 Am J Ind Med 36 415-422 Diesel exhaust and lung cancer mortality in potash mining

Schepers GWH 1963 Diseases of the chest 44 133-40 Lung disease caused by inorganic and organic dust

Selden A I *et al* 1997 Am J Ind Med 32(5) 467-77 Cancer Morbidity in workers at alumina foundries and secondary aluminium smelters.

**Selinger** B 1989 Chemistry in the market place 4<sup>th</sup> edition, HBJ

**Sigmond** G *et al* 1980 Study on the disposal of bauxite residues – final report United Nations Industrial Development Organization Budapest Hungary

Simonato L *et al* 1988 Carcinogenesis 9 1159-65 estimates of the proportion of lung cancer attributable to occupational exposure

Soskolne CL *et al* 1989 Archives of Environmental Health 44(3) 180-189 Epidemiological and Toxicologic evidence for chronic health effects and the underlying biologic mechanisms involved in sub-lethal exposures to acidic pollutants.

Sound Telegraph 2004 March 24 p 2 Multi-million dollar installation



Southwell M a 2002 The West Australian Jan 29 2002 p6 Red mud studies raise questions – monitoring was inadequate.

Southwell M b 2002 The West Australian February 19 p 3 Alcoa doctors conflict – Wagerup inquiry hears call for better industrial hygiene

Southwell M c 2002 The West Australian February 19 p Firm admits plant fallout scars cars

Southwell M d 2002 The West Australian March 9 Cancer rates higher near industry sites

Southwell M e 2002 The West Australian March 12 p 4 Expert attacks WA on emission regulations

Southwell M f 2002 The West Australian March 13 p 35 Alcoa warned by consultants – Wagerup emissions would be restricted in the US: report

Southwell M g 2002 The West Australian March 27 p15 Inquiry told of toxic leak fear – Department defends decision not to prosecute

Southwell M h 2002 The West Australian March 27 p15 Big caustic spill at Wagerup

Southwell M i 2002 The West Australian Apr 6 p Prosecute Alcoa over leaks, urges Edwards – spokesman for residents welcomes tough stand

Southwell M j 2002 The West Australian Apr 6 p Wagerup move caused cancer: woman

Southwell M k 2002 The West Australian Apr 12 p 4 Heavy industries slip net on pollution – Minister seeks to close loophole in environmental laws

Southwell M l 2002 The West Australian Residents look to class action against Alcoa

Southwell M m 2002 The West Australian May 2 p 1 Alcoa in \$M payout – Offer to sick workers

Southwell M n 2002 The West Australian May 24 p7 Alcoa told of health issues – Kwinana refinery benzene emissions five times the recommended maximum

Southwell M o 2002 The West Australian June 27 p3 Alcoa cancer rate shock – some diseases 50 per cent more common in mine and refinery workers

Southwell M p 2002 The West Australian June 27 p6 Health fears prompt Alcoa walkout

Southwell M q and Amalfi C 2002 The West Australian August 31 p12 Widows say jobs caused their loss

Southwell M r 2002 The West Australian September 2 p6 K58 cancer tie unproved – smoking may account for higher Alcoa cases, says WA health lobby

Southwell M s 2002 The West Australian September 2 p6 History of workers' health worries – leaked documents show Alcoa knew of health fears 10 years ago

Southwell M t 2002 The West Australian September 6 p28 Refinery health link raised

Southwell M u 2003 The West Australian March 15 p15 Yarloop air test shows high carcinogen compound

Southwell M v 2003 The West Australian March 31 p10 Emissions up: report

Southwell M w 2003 The West Australian May 24 p3 Alcoa misled: watchdog – emissions monitoring at Wagerup alumina refinery flawed

Southwell M x 2003 The West Australian p Alcoa noise report faces challenge

**Southwell M y** 2003 The West Australian July 1 p Alcoa loses self-policing right over dust tests

Southwell M z 2003 The Sunday Times p Future shock

Southwell M za 2004 The Beacon p Alcoa gets the soft touch

**Stackair** 1997 Test report Alcoa Oxalate kiln 15 April 1997

**Stanford R** 1982 Studies on radioactivity. Report to the Health Department of WA on the radioactivity of caustic bauxite residue.

Strutt J 2004 p Alcoa spill 5 times bigger

Strutt J 2004 p Alcoa report delayed

Summers RN et al 1996 Aust J Soil res 34 569-81 Bauxite residue (red mud) improves pasture growth on sandy soils in Western Australia

Summers RN and Pech JD 1997 Agriculture Ecosystems and Environment 64 219-32 Nutrient and metal content of water, sediment and soils amended with bauxite residue in the catchment of the peel Inlet and Harvey Estuary, Western Australia

**Tatrai E et al** Acta Morphol Acad Sci Hung. 1982 30(1):79-87. Experimental pneumoconiosis due to dusts of ore mines, I.

**Terry K W et al** 1998 Minerals and Energy Research Institute of Western Australia (MERIWA) Report No 195 Characterisation of inhaled dusts on mine sites

Toussaint LF 1985 Radiation Protection in Australia 3(4) 151-55 Background radiation in Western Australia

Trott P 2004 Alcoa investigated over second caustic spill

The Sunday Times 2004 November 28 Editorial Alcoa spills raise more health fears

The West Australian 2002 Editorial March 13 Alcoa steps in right direction

The West Australian 2002 May 1 p 11 Alcoa told to pay for cancer man – Victorian decision opens legal path for 7000 workers

The West Australian 2003 May 17 Writ Watch p Widow links sinus cancer to refinery

Thompson Y 2003 Community Weekend Courier June 20 p 1 State to probe cancer deaths

Thompson Y 2003 Community Weekend Courier December EPA report for public comment

Thompson Y 2004 Community Weekend Courier p Pollies back liquor burner protest

Thompson Y 2004 Community Weekend Courier October 1 p Spills record way below standards

Townsend MC et al 1985 Am Rev Respir Dis 132 1174-80 Pulmonary function in relation to total dust exposure at a bauxite refinery and alumina based chemical products plant

USDHHS 2005 U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. Report on Carcinogens, Eleventh Edition

US Environmental Protection Agency Integrated Risk Information System Diesel Engine Exhaust <http://www.epa.gov/iris/subst/0642.htm> Alcoa 8.38

Videnieks M 2004 The West Australian August 26 p38 Mine bosses face jail for lax safety

**Vincent** JH and Donaldson K 1990 Br J Ind Med 47 302-7 A dosimetric approach for relating the biological response of the lung to the accumulation of inhaled mineral dust

Vincent JH *et al* 1987 Ann Occup Hyg 31 375-93 accumulation of inhaled mineral dust in the lung and associated lymph nodes: implications for exposure and dose in occupational lung disease

Western Australia Legislative Council Report of the Standing Committee on Environment and public affairs in relation to the Alcoa Refinery at Wagerup Inquiry Report 11, October 2004

Williams R *et al* 2004 The West Australian p12 Sickness link to Alcoa plant

Woollard K 2002 The Sunday Times June 16 p32

**Zheng W *et al*** 1992 Int J Cancer. 52(4):557-61. A population-based case-control study of cancers of the nasal cavity and paranasal sinuses in Shanghai.

**Zheng W *et al*** 1993 Am J Epidemiol 138(11) 965-72 Risk factors for cancers of the nasal cavity and paranasal sinuses among white men in the United States